

Questions about *Evaluation* of the Biological Activity of Compounds: Techniques and Mechanism of Action Studies

Q1. An important chemical and mathematical concept was not described in the book chapter: what does *the Law of Mass Action* mean? (An ODE model of reaction rate and reactant mass)

Q2: Which quantity measures binding affinity directly: dissociation constant (K_D) or the concentration of the test compound that produces 50 percent inhibition (IC_{50})? (K_D)

Q3: In Figure 2.3, what do x- and y-axis represent in panel (A) and panel (B), respectively? (concentrations in in

x-axis; y-axis: counts per minute of radioactivity (A), percentage of binding of the labelled compound)

Q4: What is a sigmoidal curve? (A S-shaped, logistic or logit curve)

Q5: Do IC₅₀ values indicate a particular mechanism of action (MoA)? (No)

Q6: In a certain enzymatic assay, two compounds have the following pIC50 values: 7.2 (Compound A), 9.3

(Compound B). If all other conditions are held constant, what is the relationship between binding affinities of the two compounds with regard to the target? (B>A)

Q7: Why is DMSO often used in bioassays? (solvent, control)

Q8: Can you use your own language to describe what is the Hill function? (discussed in this lecture) Q9: What statistical measure is used to measure the signal-noise ratio in screening? Can you use your own language explaining it? (Z' score, how well can we separate positive controls from negative controls) Q10: Why logarithm (usually base 10) transformation is often preferred to represent quantities such as IC₅₀ and K_i? (presentation, as well as statistical mechanistics)

AMIDD Lecture 6: Structure- and ligand-based drug discovery



Dr. Jitao David Zhang, Computational Biologist

¹ Pharmaceutical Sciences, Pharma Research and Early Development, Roche Innovation Center Basel, F. Hoffmann-La Roche ² Department of Mathematics and Informatics. University of Basel Jaeger, Kathrin, Steffen Bruenle, Tobias Weinert, Wolfgang Guba, Jonas Muehle, Takuya Miyazaki, Martin Weber, et al. "<u>Structural Basis for Allosteric Ligand</u> <u>Recognition in the Human CC Chemokine Receptor 7.</u>" Cell 178, no. 5 (August 22, 2019): 1222-1230.e10.

Workflow in a typical drug-discovery program

- 1. Compound library construction;
- 2. Screening compounds with *bioassays*, or *assays*, which determine potency of a chemical by its effect on biological entities: proteins, cells, *etc*;
- 3. Hit identification and clustering;
- 4. More assays, complementary to the assays used in the screening, maybe of lower throughput but more biologically relevant;
- 5. Analysis of ligand-target interactions, for instance by getting the co-structure of both protein (primary target, and off-targets if necessary) and the hit;
- 6. *Drug design,* namely to modify the structure of the drug candidate;
- 7. Analog synthesis and testing (back to step 4);
- 8. Multidimensional Optimization (MDO), with the goal to optimize potency, selectivity, safety, bioavailability, *etc;*
- 9. Further *in vitro*, *ex vivo*, and *in vivo* testing, and preclinical development;
- 10. Entry into human (Phase 0 or phase 1 clinical trial).



Structure-based and ligand-based drug design





Target and its protein structure

QSAR= quantitative structure activity relationship; MoA= mechanism of action, or mode of action

Outline



- Representation of drug-like small molecules
- Affinity of drug to protein target
- The Michaelis-Menten model and enzymatic kinetics



Molecular descriptors are numeric values that describe chemical molecules.

Molecular descriptors enable **quantification of molecular properties**. They allow mathematical operations and statistical analysis that associate biophysical/biochemical properties with molecule structures.

Example: logP (*P* for partition) is an experimental molecular descriptor.



Lipinski's Rule of Five of small-molecule drugs

- HBD<=5: No more than 5 hydrogen-bond donors, e.g. the total number of nitrogen-hydrogen and oxygen-hydrogen bonds.
- HBA<=10: No more than 10 hydrogen-bond acceptors, e.g. all nitrogen or oxygen atoms
- MW<500: A molecular weight less than 500 Daltons, or 500 g/mol.
- logP<=5: An octanol-water partition coefficient (log P) that does not exceed 5. (10-based)



U N I B A S E I



Rules are made to be broken

drug	year approved	therapeutic area	MW	cLogP	HBD	N+O
velpatasvir	2016	HCV	883.02	2.5	4	16
venetoclax	2016	oncology	868.44	10.4	3	14
elbasvir	2016	HCV	882.0	2.6	4	16
grazoprevir	2016	HCV	766.90	-2.0	3	15
cobimetinib	2015	oncology	531.31	5.2	3	5
daclatasvir	2015	HCV	738.88	1.3	4	14
edoxaban	2015	cardiovascular	548.06	-0.9	3	11
ombitasvir	2014	HCV	894.13	1.3	4	15
paritaprevir	2014	НСУ	765.89	1.1	3	14
netupitant	2014	nausea from chemotherapy	578.59	6.8	0	5
ledipasvir	2014	HCV	889.00	0.9	4	14
ceritinib	2014	oncology	558.14	6.5	3	8

Table 1. New FDA Approvals (2014 to Present)a of Oral bRo5 Drugs

DeGoey, *et al.* 2018. "<u>Beyond the Rule of 5: Lessons</u> <u>Learned from AbbVie's Drugs and Compound Collection.</u>" Journal of Medicinal Chemistry 61 (7): 2636–51.

Molecular descriptors of increasing complexity



Extended-connectivity fingerprints (ECFPs) and Functional-class fingerprints (FCFPs) extracting and comparing (multi-)sets of subgraphs



Implemented in <u>RDKit</u> and other software. Publication and tutorials: (1) Rogers, David, and Mathew Hahn. "<u>Extended-Connectivity Fingerprints.</u>" Journal of Chemical Information and Modeling (2010). (2) Tutorial by <u>Manish Kumar</u> and (3) Tutorial by <u>Leo Klarner</u>.



Molecular similarity and similarity measures



Table 2 Formulas for the various similarity and distance metrics

Distance metric	Formula for continuous variables ^a	Formula for dichotomous variables ^a	
Manhattan distance	$D_{A, B} = \sum_{j=1}^{n} x_{jA} - x_{jB} $	$D_{AB} = a + b - 2c$	
Euclidean distance	$D_{A, B} = \left[\sum_{j=1}^{n} (x_{jA} - x_{jB})^2\right]^{1/2}$	$D_{A,B} = [a+b-2c]^{1/2}$	
Cosine coefficient	$S_{A,B} = \left[\sum_{j=1}^{n} x_{jA} x_{jB}\right] / \left[\sum_{j=1}^{n} (x_{jA})^2 \sum_{j=1}^{n} (x_{jB})^2\right]^{1/2}$	$S_{A,B} = \frac{c}{[ab]^{1/2}}$	
Dice coefficient	$S_{A,B} = \left[2\sum_{j=1}^{n} x_{jA} x_{jB}\right] / \left[\sum_{j=1}^{n} (x_{jA})^{2} + \sum_{j=1}^{n} (x_{jB})^{2}\right]$	$S_{AB} = 2c/[a+b]$	
Tanimoto coefficient	$S_{A,B} = \frac{\left[\sum_{j=1}^{n} x_{jA} x_{jB}\right]}{\left[\sum_{j=1}^{n} (x_{jA})^{2} + \sum_{j=1}^{n} (x_{jB})^{2} - \sum_{j=1}^{n} x_{jA} x_{jB}\right]}$	$S_{AB} = c/[a+b-c]$	
Soergel distance ^b	$D_{A, B} = \left[\sum_{j=1}^{n} x_{jA} - x_{jB} \right] / \left[\sum_{j=1}^{n} max(x_{jA}, x_{jB})\right]$	$D_{A,B} = 1 - \frac{c}{[a+b-c]}$	

S denotes similarities, while *D* denotes distances. The two can be converted to each other by *similarity=1/(1+distance)*. x_{jA} means the j-th feature of molecule A. a is the number of *on* bits in molecule A, b is number of *on* bits in molecule B, while c is the number of bits that are *on* in both molecules.

(Left) Maggiora, Gerald, Martin Vogt, Dagmar Stumpfe, und Jürgen Bajorath. "Molecular Similarity in Medicinal Chemistry". Journal of Medicinal Chemistry 57, Nr. 8 (24. April 2014): 3186–3204. (Right) Bajusz, Dávid, Anita Rácz, and Károly Héberger. 2015. "Why Is Tanimoto Index an Appropriate Choice for Fingerprint-Based Similarity Calculations?" Journal of Cheminformatics 7 (1): 20.





Quantitative Structure-Activity Relationships (QSARs)

QSAR is a statistical modelling of correlation between biological activity and physicochemical properties, or $\Delta \phi = f(\Delta S)$, where ϕ indicates a biological activity and S indicates a chemical structure (1868-1869).

		Target property	MD ₁	MD ₂		MD _M
G	C ₁	У ₁	x _{1,1}	х _{1,2}		х _{1,М}
s (C ₂	У ₂	x _{2,1}			
pu	C ₃	У ₃				
	C ₄	У ₄				
Ĩ						
ပိ						
	C _N	У _N	X _{N,1}	X _{N,2}		x _{N,M}

Molecular Descriptors (MD)

The basic form of a QSAR model: find a function *f* that predicts *y* from *x*, $y \sim f(x)$

An example: **The Free-Wilson analysis.** The assumption: the biological activity for a set of analogues could be described by the contributions that substituents or structural elements make to the activity of a parent structure.





QSAR models mark the early adoption of statistical modelling and machine learning in drug discovery, the fifth type of mathematical modelling

- QSAR is among the earliest subjects that used machine learning and pattern recognition in drug discovery.
- Advantages: technically easy, fast, and many models are useful as filters.
- Disadvantages: statistical models cannot capture mechanistic aspects of biochemical interactions, limited ability to debug when a model fails to work, and findings may not be generalizable.



Interpretable and Causal Models will become more important



UNI BASEL

Performance

Offline activities

UNI BASEL

- Required reading: Badillo *et al.* 2020. "<u>An Introduction to Machine Learning</u>." Clinical Pharmacology & Therapeutics.
- Read selected pages of *Computational Methods in Drug Discovery* by Sliwoski *et al.* Please submit your results to <u>the Google Form</u> by November 11th.
- Fill the <u>anonymous survey #6</u>.



Resources and backup slides



The general practice of training a supervised learning model



(Left) To assess the generalization ability of a supervised learning algorithm, data are separated into a training subset used for building the model and a test subset used to assess the generalization error (from Badillo *et al.*, 2020) (Right) Temporal validation is especially important for drug discovery, because chemical structures used in the training set may differ substantially from those that will be tested.



Molecular similarity does not equal biological similarity



Duran-Frigola, Miquel, Eduardo Pauls, Oriol Guitart-Pla, Martino Bertoni, Víctor Alcalde, David Amat, Teresa Juan-Blanco, and Patrick Aloy. 2020. "Extending the <u>Small-Molecule Similarity Principle to All Levels of Biology with the Chemical Checker</u>." Nature Biotechnology, May, 1–10.

Resources for learning about machine learning



UNI BASEL

DataWarrior: an open-source program for data visualization and analysis with chemical intelligence



DataWarrior was and still is developed at Actelion/Idorsia Pharmaceuticals Ltd.

UNI BASEI

Selected subset of functionalities

- Molecular descriptor calculation
- Similarity calculation
- Compound clustering
- Docking

Thomas Sander, Joel Freyss, Modest von Korff, Christian Rufener. DataWarrior: An Open-Source Program For Chemistry Aware Data Visualization And Analysis. J Chem Inf Model 2015, 55, 460-473, <u>doi 10.1021/ci500588j</u>

Seeing how a drug work

Work by Thomas Shafee, Shared under CC-AS-4.0, and work by Boghog. Based on PDB record 4QI9.



The protein: Dihydrofolate reductase (DHFR), which converts dihydrofolic acid into tetrahydrofolate. The process is important for cell proliferation and cell growth. DHFR is a drug target for cancer and autoimmune diseases.

The natural substrate: dihydrofolic acid (vitamin B9), in black. Dihydrofolic acid is the *natural ligand* of DHFR.

The drug: methotrexate (MTX), in green. MTX is a *synthesized ligand* of DHFR, and it is a competitive inhibitor of DHFR with regard to its natural substrate.

The binding site: where the enzyme binds its substrate and catalyses the chemical reaction, in <u>blue</u>.

22

The biophysical and biochemical views of ligand-target binding

- The **biophysical view** of binding: Both enthalpy (heat transfer) and entropy (disorder) contribute to the binding energy ($\Delta G = \Delta H T\Delta S$).
 - Binding occurs in favourable steric, *i.e.* spatial, configurations. A simplification is **the 'lock-and-the-key' model**, however, in reality enzyme undergoes changes in its shape.
 - Binding is mediated by intermolecular forces, such as electrostatic interactions (*e.g.* hydrogen bonds), Van der Waals forces (dipole interactions), π-effects (interactions of π-orbitals of a molecular system), and hydrophobic effect.
 - Binding opposes motion, and motion opposes binding: there is enthalpy/entropy compensation in ligand-substrate binding.
- The biochemical view of binding: The rate of binding is called affinity, often expressed in K_d or, for inhibitors, K_i. A closely related, and often confusing, concept is IC₅₀.





Four basic types of kinetic mechanism of inhibition, source: <u>sciencesnail.com</u>





From the law of mass action to ligand-target interaction



The biophysical (thermodynamic) view of binding affinity: enthalpy and entropy





Kinetic and thermodynamic measurements of two p38 α inhibitors. (A) The time course of SB203580 binding to immobilized mitogen activated kinase p38 α . The y-axis shows the mass change resulting from compound binding to p38 α . At t=0, a range of SB203580 concentrations were passed across the immobilized p38 α to measure net association, and then at t=50s, the compound is replaced with buffer to initiate dissociation. The table shows the association and dissociation rate constants as well as the equilibrium dissociation constants (KD(M)) for two compounds. (B) Thermodynamic analysis. Enthalpy and entropy components of binding derived from the Van't Hoff analysis are detailed in the attached table. Δ G, Δ H and T Δ S values are in kJ/mol.

For a thorough discussion about enthalpic and entropic contributions to molecular interactions, see <u>A Medicinal</u> <u>Chemist's Guide to Molecular Interactions</u> (Journal of Medicinal Chemistry 53 (14): 5061–84) by Bissantz et al.



The biochemical (kinetic) view of binding affinity: the hyperbola curve and the dissociation constant K_D



Binding assays with direct and indirect measurements. (A) A direct binding assay using I¹²⁵ labelled cyanopindolol as a β 2-adrenoceptor ligand. The curve describes a rectangular hyperbola which saturates at high ligand concentration. The ligand dissociation constant (KD) was estimated as 0.3 nM and is a measure of the ligand affinity. (B) A typical inhibition analysis using membranes expressing the human β 2-adrenoceptor and employing 0.1 nM I¹²⁵ cyanopindolol as the labeled ligand. The displacing ligand, the selective β 2-adrenoceptor antagonist ICI 118551, produces complete inhibition of the specific binding yielding an IC50 of 1 nM. From *Evaluation of the Biological Activity of Compounds: Techniques and Mechanism of Action Studies*, by Iain G. Dougall and John Unitt.

Questions: (1) how can we interpret the hyperbola curve? (2) if f(x) is a function with the form of Ax/(k+x), what will be the form of function g(f(x)) where g(x)=Bx/(k'+x)? What implications does this have?



Modelling enzyme kinetics with the Michaelis-Menten model





The dose-response curve and IC50: The Hill function and *in vitro* pharmacology

- The Hill function is one of the mostly useful non-linear functions to model biological systems.
- In its general form, H_{max} indicates the maximal value to which the function is asymptotic, n is the shape parameter (known as the Hill's coefficient), and k is the reflection point, often abbreviated as XC₅₀ (X=I, E, C, ...), the half-saturation constant.
- The Michaelis-Menten model is a special case of the Hill function (*n*=1).



Suppose it is an antiviral drug, compared with curve B, what does curve A, C, and D suggest?



Theoretical and practical considerations about the Hill function

- The Hill function is often used to model either *target occupancy* or *tissue response*. In pharmacology, it is often used to model the tissue response.
- The Hill function can be approximated by a step function when *n* goes towards infinity (top panel). This can be seen as one of the theoretical foundations of Boolean network modelling.
- The Hill function can be deduced from statistical mechanics of binding, a particle modelling approach. See for instance an <u>article on Biophysics Wiki by Andreas Piehler</u> for details.
- Dose-response data may look quite different from the ideal curve (bottom panel). By using a Bayesian inference approach (setting priors), it is possible to perform inference even with ill-looking data.

The Bayesian inference approach versus the non-Bayesian Marquardt-Levenberg algorithm for non-linear regression fitting (other options are gradient descent and Gauss-Newton methods). 4P: four parameter model; 2P: two parameter model (IC50 and *n*). Figure 2 from Labelle, Caroline, Anne Marinier, and Sébastien Lemieux. 2019. "Enhancing the Drug Discovery Process: Bayesian Inference for the Analysis and Comparison of Dose–Response Experiments." Bioinformatics 35 (14): i464–73.







Four classical classes of mathematical models

Compartment models

$$rac{d[LR]}{dt}=k_1[L][R]-k_2[L]$$

Kinetics of ligand-target

 $\frac{dx}{dt} = \alpha x - \beta xy,$ $\frac{dy}{dt} = -\gamma y + \delta xy,$ get The Lotka-Volterra equations modelling predator-prey relationships.

$$\begin{split} \frac{dS}{dt} &= -\frac{\beta IS}{N}, \\ \frac{dI}{dt} &= \frac{\beta IS}{N} - \gamma I, \\ \frac{dR}{dt} &= \gamma I \end{split}$$

The SIR (S=susceptible, I=infectious, R=removed) model of epidemiology

Particle models



A Study on Socio-spatial Segregation Models Based on Multi-agent Systems by Quadros *et al.* (2012). 10.1109/BWSS.2012.14.

Transport models



McGinty, Sean, and Giuseppe Pontrelli. 2015. "<u>A General Model of</u> <u>Coupled Drug Release and Tissue</u> <u>Absorption for Drug Delivery</u> <u>Devices</u>." Journal of Controlled Release 217 (November): 327–36.

Finite state models



A finite-state Markov chain modelling DNA sequences

Summary and Q&A





Ligand-based approaches



Today we learned **molecular modelling techniques** and **ligand-target interaction:**

- (A) 3D protein structure-based approaches. An example with docking can be found in the backup slides.
- (B) Ligand-based approaches (similarity search). Another example of amphiphilicity can be found in the backup slides.

Zhang, Jitao David, Lisa Sach-Peltason, Christian Kramer, Ken Wang, and Martin Ebeling. 2020. "<u>Multiscale Modelling of Drug</u> <u>Mechanism and Safety</u>." *Drug Discovery Today* 25 (3): 519–34.

Summary of basic concepts



- Ligand: the binding partner of a macromolecule (often proteins), for instance other proteins (in case of protein-protein inaction), substrates and allosteric modulators (in case of enzymes). Many drugs are ligands of proteins.
- **Binding:** the formation of interactions between a protein and its ligand. In drug discovery, we encounter more often ٠ transient and non-covalent interaction (i.e. no sharing of electrons between atoms), but there are drugs form reversible or irreversible covalent bonds.
- Non-covalent interaction: electromagnetic interactions between molecules or within a molecule without forming a chemical bond, *i.e.* no sharing of electrons between atoms. Non-covalent interactions are classified into four categories: electrostatic, van der Waals forces, hydrophobic effects, and π -effects. See <u>Wikipedia</u> for more details of these interactions.
- **Conformational change**: ligand binding often triggers a change in the shape of the protein, which alters its cellular function
- Agonist versus antagonist: an agonist activates the function of its target by binding, and an antagonist blocks the action of the target by binding.
- Active site versus allosteric site: active site is where the enzyme-substrate interaction happens, example: at the active site oxygen binds to heme, and CO can compete with oxygen for heme binding. Allosteric site (i.e. regulatory site) is any other site than the active site where a ligand can bind to modulate the protein function.



More about molecular interactions and drug design

- Molecular interactions for drug discovery
 - Bissantz, Caterina, Bernd Kuhn, and Martin Stahl. "A Medicinal Chemist's Guide to Molecular Interactions." Journal of Medicinal Chemistry 53, no. 14 (July 22, 2010): 5061–84.
 <u>https://doi.org/10.1021/jm100112j</u>. A comprehensive introduction to common types of interactions, their applications, and caveats of blindly following rules in drug design.
 - Persch, Elke, Oliver Dumele, and François Diederich. "Molecular Recognition in Chemical and Biological Systems." Angewandte Chemie International Edition 54, no. 11 (2015): 3290–3327.
 <u>https://doi.org/10.1002/anie.201408487</u>. A comprehensive introduction to molecular recognition.
- How drug design help with drug discovery: ten real-life stories
 - Kuhn, Bernd, Wolfgang Guba, Jérôme Hert, David Banner, Caterina Bissantz, Simona Ceccarelli, Wolfgang Haap, et al. "A Real-World Perspective on Molecular Design." Journal of Medicinal Chemistry 59, no. 9 (May 12, 2016): 4087–4102. <u>https://doi.org/10.1021/acs.jmedchem.5b01875</u>. The common themes summarized in the Conclusion are helpful in my opinion for any scientist working in quantitative aspects of drug discovery: (1) value of qualitative statements, (2) shaping chemical space, (3) the principle of parsimony, (4) annotation is half the battle, and (5) staying close to experiment.

The principle of molecular docking, a case study of structure-based drug design

Docking is like a discotheque: it is all about posing and scoring – Roger Sayle (*NextMove* Software Limited)

Three basic methods to represent target and ligand structures *in silico*

- *Atomic*: used in conjunction with a potential energy function, computational complexity high
- Surface: often used in protein-protein docking
- *Grid representation*: the basic idea is that to store information about the receptor's energetic contributions on grid points so that it only needs to be read during ligand scoring.

In the most basic form, grid points store two types of potentials: **electrostatic** and **van der Waals forces**, for instance using Coulombic interactions and L-J 12-6 function.

 $E_{coul}(r) = \sum_{i=1}^{N_A} \sum_{j=1}^{N_B} \frac{q_i q_j}{4\pi\varepsilon_0 r_{ij}}$

$$E_{vdW}(r) = \sum_{j=1}^{N} \sum_{i=1}^{N} 4\varepsilon \left[\left(\frac{\sigma_{ij}}{r_{ij}} \right)^{12} - \left(\frac{\sigma_{ij}}{r_{ij}} \right)^{6} \right]$$

- ε is the **well depth** of the potential
- σ is the collision diameter of the respective atoms *i* and *j*.



Kitchen, Douglas B., Hélène Decornez, John R. Furr, und Jürgen Bajorath. "Docking and Scoring in Virtual Screening for Drug Discovery: Methods and Applications". *Nature Reviews Drug Discovery* 3, Nr. 11 (November 2004): 935–49. <u>https://doi.org/10.1038/nrd1549</u>.

Coulombic interactions (electrostatic interactions between electric charges)

> Lennard–Jones 12–6 function

(intermolecular interactions without charge)





Posing: dealing with flexibility of ligand and of protein



Chen, Yu-Chian. "Beware of docking!" *Trends in Pharmacological Sciences* 36, Nr. 2 (1. Februar 2015): 78–95. https://doi.org/10.1016/j.tips.2014.12.001.





Methods to deal with ligand and protein flexibility

- Systematic search
- Random search, such as Monte-Carlo and genetic algorithms
- Simulation methods, such as molecular dynamics

UNI BASEL

Types of scoring functions





- Empirical scoring functions estimate the binding affinity of a complex by summing up the important energetic factors for protein-ligand **binding**, such as hydrogen bonds, hydrophobic effects, steric clashes, etc. It relies on training set and regression analysis.
- Knowledge-based scoring functions derive the desired pairwise potentials from three-dimensional structures of a large set of protein-ligand complexes based on the inverse Boltzmann distribution. It is assumed that the frequency of different atom pairs in different distances is related to the interaction of two atoms and converts the frequency into the distance-dependent potential of mean force.
- Machine learning-based scoring functions are usually used for rescoring to improve the initial docking.

Li, Jin, Ailing Fu, und Le Zhang. "An Overview of Scoring Functions Used for Protein-Ligand Interactions in Molecular Docking". Interdisciplinary Sciences: Computational Life Sciences 11, Nr. 2 (1. Juni 2019): 320-28. https://doi.org/10.1007/s12539-019-00327-w.

UNI BASEI

Interested in learning more about molecular modelling?



PROTOCOL

Computational protein-ligand docking and virtual drug screening with the AutoDock suite

Stefano Forli, Ruth Huey, Michael E Pique, Michel F Sanner, David S Goodsell & Arthur J Olson

Department of Integrative Structural and Computational Biology, The Scripps Research Institute, La Jolla, California, USA. Correspondence should be addressed to A.J.O. (olson@scripps.edu).

Published online 14 April 2016; doi:10.1038/nprot.2016.051

Computational docking can be used to predict bound conformations and free energies of binding for small-molecule ligands to macromolecular targets. Docking is widely used for the study of biomolecular interactions and mechanisms, and it is applied to structure-based drug design. The methods are fast enough to allow virtual screening of ligand libraries containing tens of thousands of compounds. This protocol covers the docking and virtual screening methods provided by the AutoDock suite of programs, including a basic docking of a drug molecule with an anticancer target, a virtual screen of this target with a small ligand library, docking with selective receptor flexibility, active site prediction and docking with explicit hydration. The entire protocol will require ~5 h.

- Try docking yourself by following this protocol: Forli, Stefano, Ruth Huey, Michael E. Pique, Michel F. Sanner, David S. Goodsell, und Arthur J. Olson. "Computational Protein–Ligand Docking and Virtual Drug Screening with the AutoDock Suite". *Nature Protocols* 11, Nr. 5 (Mai 2016): 905–19. <u>https://doi.org/10.1038/nprot.2016.051</u>.
- In-depth reading: Sliwoski, Gregory, Sandeepkumar Kothiwale, Jens Meiler, und Edward W. Lowe. "Computational Methods in Drug Discovery". *Pharmacological Reviews* 66, Nr. 1 (1. Januar 2014): 334–95. <u>https://doi.org/10.1124/pr.112.007336</u>.
- A more advanced talk by Arthur Olson can be found <u>here</u>, Workshop on the Mathematics of Drug Design/Discovery, June 4 8, 2018, The Fields Institute. Courses available at the University of Basel and beyond.
- Binding predicted by docking should always be challenged and verified by experimental testing! Docking scores seldomly correlate with binding affinity.

More about the the Free-Wilson analysis

- <u>A Mathematical Contribution to Structure-Activity Studies</u> by Spencer M. Free and James W. Wilson, Journal of Medicinal Chemistry, 1964, and reviewed by <u>Kubinyi</u>, 1988.
- A Python implementation on <u>GitHub</u>, and a <u>blog post</u> going through examples, is shared by Pat Walters.
- Free-Wilson nonadditivity is a research topic, for instance see Cramer et al., 2015
- Source of the example shown in the lecture: QSAR of the <u>ACCVIP</u> project (The Australian Computational Chemistry via the Internet Project)

UNI BASEI

Drug-induced phospholipidosis is correlated with amphiphilicity



- Phospholipidosis is a lysosomal storage disorder characterized by the excess accumulation of phospholipids in tissues.
- Drug-induced phospholipidosis is caused by cationic amphiphilic drugs and some cationic hydrophilic drugs.
- Clinical pharmacokinetic characteristics of drug-induced phospholipidosis include (1) very long terminal half lives, (2) high volume of distribution, (3) tissue accumulation upon frequent dosing, and (4) deficit in drug metabolism.

Fischer *et al.* (Chimia 2000) discovered that it is possible to predict the amphiphilicity property of druglike molecules by calculating the amphiphilic moment using a simple equation.



Lüllmann *et al.*, Drug Induced Phospholipidosis, *Crit. Rev. Toxicol. 4, 185, 1975*



Anderson and Borlak, Drug-Induced Phospholipidosis,. *FEBS Letters* 580, Nr. 23 (2006): 5533–40.



 \vec{A} : Caculated amphiphilic moment

d: distance between the center of gravity of the charged part of a molecule and the hydrophobic/hydrophilic remnant of the molecule

 $\vec{\alpha}_i$: the hydrophobic/hydrophilic contribution of atom/fragment *i*



In silico prediction of amphiphilicity

Development of CAFCA (<u>CA</u>lculated <u>Free</u> energy of amphiphilicity of small <u>C</u>harged <u>A</u>mphiphiles)



Iterative model building, experimentation, and model refining led to the predictive tool CAFCA

Validation of in silico phospholipidosis prediction



Model Validation from 1999-2004



Calculated Basic pKa

Plot of amphiphilicity ($\Delta\Delta G_{AM}$) versus calculated basic pK_a for the training set of 24 compounds. The red area defines the region where a positive PLD response is expected, and the green area defines where a negative response is expected according to the tool.

in vitro/	in silico/	Exp. PC/	In silico/	n=36
in vivo	in vivo	in vivo	in vitro	
94%	81%	89%	89%	

in	n=422		
Accuracy [(TP+TN)/ (P+N)]	Sensitivity [True Positive Rate]	Specificity [True Negative Rate]	Precision [TP/(TP+FP)]
86%	80%	90%	84%

Fischer et al., J. Med. Chem, 55 (1), 2012

We gained mechanistic insights of phospholipidosis induction by cationic amphiphilic drugs with the model

Phospholipidosis: lessons learned (and lessons not yet learned)

- Cationic amphiphilic properties of a molecule is an early marker for safety in drug discovery and early development.
 - Phospholipidosis in dose range finding studies
 - Cardiac ion channel interactions (hERG, natrium channel, ...)
 - Receptor binding promiscuity
 - P-gp inhibition
 - Mitochondrial toxicity in case of safety relevant findings, e.g. in dose range finding studies
- Extreme basic amphiphilic properties should be avoided because of a higher risk of PLD, QT-prolongation, mitochondrial toxicity. However, basic compounds with moderate amphiphilic properties are still a preferred scaffold for many therapeutic areas (especially CNS).
- Generally, some safety liabilities, despite complex underlying biological and chemical mechanisms, can be predicted by molecular modelling well, sometimes with surprisingly elegant models!



Fig. 1. Representative examples of CADs that are identified in SARS-CoV-2 drug repurposing screens.

Tummino, Tia A., Veronica V. Rezelj, Benoit Fischer, Audrey Fischer, Matthew J. O'Meara, Blandine Monel, Thomas Vallet, et al. "Drug-Induced Phospholipidosis Confounds Drug Repurposing for SARS-CoV-2." Science 373, no. 6554 (July 30, 2021): 541–47. <u>https://doi.org/10.1126/science.abi4708</u>.



Resources about the mathematics underlying molecular structure determination

- Mathematical and physical foundations
 - Recommended reading: <u>Mathematical techniques used in</u> <u>biophysics</u>
 - <u>Background on imaging physics</u> at xrayphysics.com
 - <u>Physics for life-science students</u> at U Maryland
- X-ray diffraction by electrons
 - An <u>AMS Feature Column</u> by Tony Phillips
 - Stanford open course <u>Fourier transform and its applications</u>
- Nuclear Magnetic Resonance (NMR)
 - <u>A beautiful video tutorial</u> about the principles of magnetic resonance imaging (MRI), which is a variant of NMR
- Cryo-electron microscopy (CryoEM)
 - <u>A three-minute introduction to CryoEM</u>
 - <u>Nobel Prize Talk by Joachim Frank</u>
 - <u>Talk on Mathematics of CryoEM</u>, by Prof Amit Singer, with a manuscript available at arXiv: <u>https://arxiv.org/abs/1803.06714</u>



Swiss Light Source, the synchrotron at the Paul Scherrer Institute (PSI), copyright of PSI





Adapted from Bushberg JT, <u>The</u> <u>Essential Physics</u> <u>of Medical</u> <u>Imaging</u>: Lippincott Williams & Wilkins; 2002

N M NEXUS/Physics Physics for life-science students